

On nonstandard Padé approximants suitable for effective properties of two-phase composite materials

Miao-jung Y. Ou

UT Joint Institute for Computational Sciences, Oak Ridge National Laboratory,
Oak Ridge, TN 37831, USA*
mou@utk.edu

February 25, 2010

Abstract

This paper investigates existence of the nonstandard Padé approximants introduced by Cherkhev and Zhang in [2] for approximating the spectral function of composites from effective properties at different frequencies. The spectral functions contain microstructure information. Since this reconstruction problem is ill-posed [9], the well-performed Padé approach is noteworthy and deserves further investigations. In this paper, we validate the assumption that the effective dielectric component of interest of all binary composites can be approximated by Padé approximants whose denominator has nonzero power one term. We refer to this as the nonstandard Padé approximant, in contrast to the standard approximants whose denominator have nonzero constant terms. For composites whose spectral function assumes infinitely many different values such as the checkerboard microstructure, the proof is carried by using classic results for Stieltjes functions. For those with spectral functions having only finitely many different values, we prove the results by utilizing a special product decomposition of the coefficient matrix of the Padé system. The results in this paper can be considered as an extension of the Padé theory for Stieltjes functions whose spectral function take infinitely many different values to those taking only finitely many values. In the literature, the latter is usually excluded from the definition of Stieltjes functions because they correspond to rational functions, hence convergence of their Padé approximants is trivial. However, from an inverse problem point of view, our main concern is the existence of the nonstandard Padé approximants, rather than their convergence. The results in this paper provide a mathematical foundation for applying the Padé approach for reconstructing the spectral functions of composites whose microstructure is not *a priori* known.

Keywords: Effective properties of composites; Microstructure; Spectral functions; Nonstandard Padé approximants; Stieltjes functions; Inverse homogenization.
AMS Subject Classification: 30E10

*Building 6012, MS 6367, 1 Bethel Valley Road, Oak Ridge, TN 37831, USA

1 Introduction

In the seminal paper by Golden and Papanicolaou[1], components of the effective dielectric matrix of a two-phase composite with statistically homogeneous microstructure and isotropic constituents are represented as Stieltjes integrals with positive Borel measure supported in $[0, 1]$. We refer to this integral representation formula as IRF throughout the paper. In the IRF, the contrast of the constituents, represented as the ratio (denoted by h) of the two dielectric constants, stays in the integrand while the information of microstructure is contained in the positive Borel measure. To be more specific, it is shown in [1] that for $s := \frac{1}{1-h}$ outside $[0, 1]$ on the complex plane, the (i, k) -component of the effective dielectric tensor, $i, k = 1, 2, 3$ of a dielectric composite, denoted by ϵ_{ik}^* can be represented as

$$\epsilon_{ik}^* = \epsilon_1 \left(\delta_{ik} - \int_0^1 \frac{1}{s-z} d\mu_{ik}(z) \right)$$

where $h := \frac{\epsilon_2}{\epsilon_1}$, δ_{ik} is Kronecker delta, and $d\mu_{ik}(z)$ is a positive Borel measure. Throughout the paper, we adapt the notation $:=$ and $=$: such that

$$A := B \Leftrightarrow A \text{ is defined by } B$$

$$A =: B \Leftrightarrow B \text{ is defined by } A$$

Following the convention in literature, we introduce $F(s)$ function

$$F_{ik}(s) := \delta_{ik} - \frac{\epsilon_{ik}^*}{\epsilon_1} = \int_0^1 \frac{1}{s-z} d\mu_{ik}(z) \quad (1)$$

Note that h is complex-valued for lossy materials. Applying geometric series expansion of $\frac{1}{s-z}$ around $s = \infty$ (corresponding to a homogeneous material), (1) becomes

$$F_{ik}(s) = \sum_{m=0}^{\infty} \frac{c_m}{s^{m+1}}$$

with c_m being the m -th moment of the measure, i.e. $c_m := \int_0^1 z^m d\mu_{ik}(z)$.

It is also shown in [1] that c_m , $m = 0, 1, 2, \dots$, is related to the $(m+1)$ -point correlation function of the characteristic function of the region occupied by one of the constituents. This link has been exploited for estimating the bounds on effective dielectric tensor by using information on m -point correlation functions such as volume fraction ($m = 1$) and two-point correlation function. This approach of estimating bounds relies on the observation from (1) that the $F_{ik}(s)$ is an analytic function for $s \in \mathbb{C} \setminus [0, 1]$, where $\mathbb{C}$ is the complex plane. This provides a mathematical scheme for retrieving the moments c_m when the dielectric constant of the one of the constituents varies with frequency. In [3], it was demonstrated analytically and numerically that the construction of moments of the measure from data of ϵ_{ik}^* at various frequencies is a stable process, unlike the ill-posed process of reconstructing the measure itself. In that paper, the reconstruction of moments

was carried out by solving a truncated linear system if all the data points s_k are outside the unit circle and by a Gaussian quadrature method. In all the tested numerical examples, the first 5-10 moments can be reconstructed with high accuracy by both methods. These moments contain the microstructural information and the resulting truncated series also provide a very accurate way for calculating the effective dielectric constant for all frequencies(extrapolation) [3]. In [2], instead of reconstructing the coefficients of the truncated Stieltjes series, the coefficients of the nonstandard Padé approximants of the series in (1) was reconstructed from data of ϵ_{ik}^* at different frequencies. This method is able to reconstruct the first 10-15 moments with high accuracy and the resulting Padé apprximant provides a very accurate way for calculating the effective dielectric constant for all frequencies(extrapolation). It is worth noticing that all the real-valued poles of the resulting Padé approximant lie along the graph of the true spectral functions in all the tested numerical examples. To be more specific, in [2], Padé approximants for the Stieltjes integral in (1) with the following form:

$$F(s) \simeq \frac{a(s)}{b(s)} = \frac{a_0 + a_1s + a_2s^2 + \cdots + a_ps^p}{b_0 + b_1s + b_2s^2 + \cdots + b_qs^q}, \quad b_1 \neq 0. \quad (2)$$

was used for recovering the moments. They assume $F(s)$ has at least one pole and normalize b_1 to be 1, unlike the standard Padé approximants for which b_0 is normalized to 1. We refer to (2) as **nonstandard Padé apprximants**. The main reason prompting the choice of $b_1 \neq 0$ is because for a checkerboard microstructure, the spectral function has a pole at $s = 0$. For ease of notation, we will replace ϵ_{ik}^* with ϵ^* hereafter. μ is assumed to take at least two different values because otherwise F will be identically zero as implied by (1). For N different frequencies $\omega_1, \dots, \omega_N$, there are N data pairs (s_k, d_k) , where $s_k = \frac{1}{1 - \epsilon_2(\omega_k)/\epsilon_1(\omega_k)}$ and $d_k = \epsilon^*(s_k)$, $k = 1, \dots, N$. The following linear system of equations for $a_0, \dots, a_p, b_0, b_2, \dots, b_q$,

$$d_k = \frac{a_0 + a_1s_k + a_2s_k^2 + \cdots + a_ps_k^p}{b_0 + s_k + b_2s_k^2 + \cdots + b_qs_k^q}, \quad p + q + 1 \leq N \quad (3)$$

was solved by formulating it as a least-square minimization problem subject to the constraints

$$0 \leq A_n < 1, \quad 0 \leq s_n < 1, \quad 0 < \sum_n A_n < 1 \quad (4)$$

for the residues A_n and poles s_n defined by the partial fraction decomposition of the nonstandard Padé approximant

$$\frac{a_0 + a_1s + a_2s^2 + \cdots + a_ps^p}{b_0 + b_1s + b_2s^2 + \cdots + b_qs^q} =: \sum_{n=1}^q \frac{A_n}{s - s_n}$$

The moments were then computed by applying partial fraction decomposition to the Padé approximants and by correctly combining the poles and residues. See [2, 4] for details.

In [5][6][7], the fact that a Stieltjes function can be bounded up and below by standard Padé approximants was used for deriving bounds on the effective dielectric constants.

The main purpose of this paper is twofold. The first is to show that $F(s)$ can always be approximated by a Padé approximant of the form in (2) for $s \in \mathbb{C} \setminus [0, 1]$, i.e. to show that we can always normalize b_1 to be 1 and to show this special form of Padé approximant has the accuracy-through-order property that will be defined in the next section. The second is to justify the constraints shown in (4).

This paper is organized as follows. We first define an auxiliary function $f(\xi)$ and prove that it has accuracy-through-order standard Padé approximants in Section 2.1. In the same section, we also prove properties of the poles of the Padé approximants. Using these results, we show in Section 3 that $F(s)$ always has accurate-through-order Padé approximant of the form as in (2) for appropriate p and q . The major technique in the proof is factorizing the coefficient matrix of the system of Padé equations into products of Vandermonde matrices and a diagonal matrix. In Section 4, we discuss the relevance of this work in materials science for binary composites.

2 Existence of Padé approximants

Since the representation formula for $F(s)$ is a Stieltjes integral with non-decreasing distribution function μ and is analytic for $s \in \mathbb{C} \setminus [0, 1]$, we may identify it with a Stieltjes function $f(\xi)$ [8] by rewriting $F(s)$ in terms of a new variable $\xi := -1/s$

$$F(s) =: G(\xi) = -\xi \int_0^1 \frac{d\mu(z)}{1 + z\xi} =: -\xi \cdot f(\xi) \quad (5)$$

We refer to $f(\xi)$ as the **auxiliary function** and would like to remark that the function $\mu(z)$ can take finitely many or infinitely many different values, i.e. the definition of Stieltjes function we use here does not include the restriction on the number of different values $\mu(z)$ takes. Note that the Stieltjes function $f(\xi)$ is analytic for $\xi \in \mathbb{C} \setminus (-\infty, -1]$ and has a power series expansion valid in $|\xi| < 1$, which reads

$$f(\xi) = \sum_{n=0}^{\infty} \mu_n (-\xi)^n. \quad (6)$$

For later use, we also include the expansion of G in the same region on the ξ -plane

$$G(\xi) = \sum_{n=0}^{\infty} \mu_n (-\xi)^{n+1}. \quad (7)$$

Here we use $-\xi$ because it is easier to work with the positive coefficients $\{\mu_n\}$, which can be easily verified as the moments of the Borel measure μ in (5), $n = 0, 1, 2, \dots$.

2.1 Existence of standard Padé approximants for the auxiliary function $f(\xi)$

Recall the standard $[L/M]$ Padé approximant for $f(\xi)$ in (6) is a rational function of the following form

$$[L/M](-\xi) := \frac{a_0 + a_1(-\xi) + \cdots + a_L(-\xi)^L}{b_0 + b_1(-\xi) + \cdots + b_M(-\xi)^M}$$

such that b_0 can be normalized to be 1 and the rest of the coefficients satisfy the system of Padé equations [8]

$$\begin{bmatrix} \mu_{L-M+1} & \mu_{L-M+2} & \cdots & \mu_L \\ \mu_{L-M+2} & \mu_{L-M+3} & \cdots & \mu_{L+1} \\ \vdots & \vdots & \ddots & \vdots \\ \mu_L & \mu_{L+1} & \cdots & \mu_{L+M-1} \end{bmatrix} \begin{bmatrix} b_M \\ b_{M-1} \\ \vdots \\ b_1 \end{bmatrix} = - \begin{bmatrix} \mu_{L+1} \\ \mu_{L+2} \\ \vdots \\ \mu_{L+M} \end{bmatrix} \quad (8)$$

$$\begin{cases} a_0 = \mu_0 \\ a_1 = \mu_1 + b_1 \mu_0 \\ a_2 = \mu_2 + b_1 \mu_1 + b_2 \mu_0 \\ \vdots \\ a_L = \mu_L + \sum_{k=1}^{\min(L,M)} b_k \mu_{L-k} \end{cases} \quad (9)$$

For consistency, we define $\mu_j = 0$ for $j < 0$. The $[L/M]$ Padé approximant is said to have the accuracy-through-order property if it satisfies

$$f(\xi) - [L/M] = O(\xi^{L+M+1}). \quad (10)$$

for $|\xi|$ within radius of convergence. We would like to remark that for any given function $f(z)$, polynomials $p^{[L/M]}(z)$, $q^{[L/M]}(z)$ of degrees L , M , respectively, can always be found so that

$$q^{[L/M]}(z)f(z) - p^{[L/M]}(z) = O(z^{L+M+1}) \quad (11)$$

However, it is well known that (11) does not necessarily imply $p^{[L/M]}(z)/q^{[L/M]}(z) = f(z) + O(z^{L+M+1})$ [8]. Following Baker, we say that the Padé approximant does not exist if the accuracy-through-order requirement is not satisfied. On the other hand, it can be shown that [8](p.7) the

rational function $\frac{P^{[L/M]}(-\xi)}{Q^{[L/M]}(-\xi)}$ with $P^{[L/M]}$ and $Q^{[L/M]}$ defined as

$$P^{[L/M]}(-\xi) = \begin{vmatrix} \mu_{L-M+1} & \mu_{L-M+2} & \cdots & \mu_{L+1} \\ \mu_{L-M+2} & \mu_{L-M+3} & \cdots & \mu_{L+2} \\ \vdots & \vdots & \ddots & \vdots \\ \mu_{L-1} & \mu_L & \cdots & \mu_{L+M-1} \\ \mu_L & \mu_{L+1} & \cdots & \mu_{L+M} \\ \sum_{i=0}^{L-M} \mu_i(-\xi)^{M+i} & \sum_{i=0}^{L-M+1} \mu_i(-\xi)^{M+i-1} & \cdots & \sum_{i=0}^L \mu_i(-\xi)^i \end{vmatrix} \quad (12)$$

$$Q^{[L/M]}(-\xi) = \begin{vmatrix} \mu_{L-M+1} & \mu_{L-M+2} & \cdots & \mu_L & \mu_{L+1} \\ \mu_{L-M+2} & \mu_{L-M+3} & \cdots & \mu_{L+1} & \mu_{L+2} \\ \vdots & \vdots & \ddots & \vdots & \vdots \\ \mu_{L-1} & \mu_L & \cdots & \mu_{L+M-2} & \mu_{L+M-1} \\ \mu_L & \mu_{L+1} & \cdots & \mu_{L+M-1} & \mu_{L+M} \\ (-\xi)^M & (-\xi)^{M-1} & \cdots & (-\xi) & 1 \end{vmatrix} \quad (13)$$

is an accuracy-through-order Padé approximant of $f(\xi)$ provided $Q^{[L/M]}(0) \neq 0$. Note that $Q^{[L/M]}(0)$ (Hankel determinant) is exactly the determinant of the coefficient matrix of (8). Hence the condition $Q^{[L/M]}(0) \neq 0$ guarantees the accuracy-through-order property of the $[L/M]$ approximant [8](p.21). If $Q^{[L/M]}(0) = 0$, (8) can have infinitely many solutions or fail to have any solution, i.e. the $[L/M]$ Padé approximant may not exist. We would like to remark that $Q^{[L/M]}(0) = 0$ does not imply non-existence of accuracy-through-order Padé approximants. For example, for a rational function with degree p numerator and degree q denominator, it is known that $Q^{[L/M]}(0) = 0$ if $L > p$ and $M > q$ but obviously, the Padé approximants are the rational function itself for these cases.

Because the numerical algorithm in [2] is based on the reconstruction of the Padé approximant of $F(s)$, rather than $F(s)$ itself, it is necessary to prove the existence of Padé approximant for $G(\xi)$ defined in (5). To achieve this, we first show the existence of standard Padé approximants for the Stieltjes function $f(\xi)$ in the following theorem.

Theorem 2.1. *The $[L/M]$ Padé approximant for $f(\xi)$ exists for all $L - M + 1 \geq 0$, $L \neq -1$ and $M \geq 0$ such that $b_0 = 1$.*

Proof. We first consider the case where the non-decreasing function $\mu(z)$ in (5) takes infinitely many different values. Using Theorem 5.2.1 in [8], which states that $Q^{[L/M]}(0) \neq 0$ for all $L - M + 1 \geq 0$ and $M \geq 1$, the existence of $[L/M]$ approximants follows directly. For $M = 0$, $[L/0]$ clearly exists because $[L/0] = \sum_{j=0}^L \mu_j(-\xi)^j$.

For the case in which $\mu(z)$ takes $n + 1$ different values for some non-negative finite integer n , we prove the theorem as follows. Let $z_1, z_2, \dots, z_n$ be the n different points on $[0, 1]$ such that $0 \leq z_1 < z_2 < z_3 < \dots < z_n$ and $\mu(z)$ changes values only at these points, i.e. $d\mu(z)$ is a finite sum of Dirac measure sitting at z_i with strength λ_i .

Case 1: $n = 0$ If $n = 0$, then $f(\xi) \equiv 0$ so $[L/M] = 0$ for all $L - M + 1 \geq 0$ and $M \geq 0$.

Case 2: $n = 1$ If $z_1 = 0$, i.e. $f(\xi)$ takes two different values and the value change occurs at z_1 by the amount of λ_1 , then $f(\xi) \equiv \lambda_1 > 0$. Clearly, the Padé approximant exists. If $z_1 > 0$, $f(\xi) = \frac{\lambda}{1+z_1\xi}$ so $[L/M] = \frac{\lambda}{1+z_1\xi}$ for all $L - M + 1 \geq 0$ and $M \geq 1$ by Theorem 1.4.4 in [8] and $[L/0]$ clearly exists because $[L/0] = \sum_{j=0}^L \mu_j(-\xi)^j$.

Case 3: $n \geq 2$ Denote the amount of value change at z_i by $\lambda_i > 0$, $i = 1, 2, \dots, n$. With this, $f(\xi)$ can be written as

$$f(\xi) = \sum_{i=1}^n \frac{\lambda_i}{1 + z_i \xi}$$

Suppose $z_1 \neq 0$, i.e. $z_i, i = 1 \sim n$ are all non-zero. Then $f(\xi)$ is a rational function with degree $n - 1$ numerator and degree n denominator. By the characterization theorem of rational function (Theorem 1.4.4) in [8], we know that $[L/M](-\xi) = f(\xi)$ if $L \geq n - 1$ and $M \geq n$, i.e. the Padé approximant of a rational function is itself if both L and M are large enough. Also, the constant term of the denominator is obviously non-zero. This takes care of those $[L/M]$ such that $L - M + 1 \geq 0$ and $M \geq n$.

To complete the proof for $z_1 \neq 0$, we consider $[L/M]$ such that $L - M + 1 \geq 0$ and $1 \leq M < n$. For this case, we exploit the structure of moments $\mu_0 = \sum_{i=1}^n \lambda_i$; $\mu_k = \sum_{i=1}^n \lambda_i z_i^k$, $k = 1, 2, 3 \dots$ to rewrite the coefficient matrix of (8) as

$$\begin{bmatrix} \sum_{i=1}^n \lambda_i z_i^{L-M+1} & \sum_{i=1}^n \lambda_i z_i^{L-M+2} & \dots & \sum_{i=1}^n \lambda_i z_i^L \\ \sum_{i=1}^n \lambda_i z_i^{L-M+2} & \sum_{i=1}^n \lambda_i z_i^{L-M+3} & \dots & \sum_{i=1}^n \lambda_i z_i^{L+1} \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{i=1}^n \lambda_i z_i^L & \sum_{i=1}^n \lambda_i z_i^{L+1} & \dots & \sum_{i=1}^n \lambda_i z_i^{L+M-1} \end{bmatrix} = \mathbf{V}^T \mathbf{\Lambda} \mathbf{V}, \quad (14)$$

where

$$\mathbf{V} := \begin{bmatrix} 1 & z_1 & z_1^2 & \cdots & z_1^{M-1} \\ 1 & z_2 & z_2^2 & \cdots & z_2^{M-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & z_M & z_M^2 & \cdots & z_M^{M-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & z_n & z_n^2 & \cdots & z_n^{M-1} \end{bmatrix}_{n \times M} \quad (15)$$

$$\mathbf{\Lambda} := \begin{bmatrix} \lambda_1 z_1^{L-M+1} & 0 & 0 & \cdots & 0 \\ 0 & \lambda_2 z_2^{L-M+1} & 0 & \cdots & 0 \\ 0 & 0 & \lambda_3 z_3^{L-M+1} & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda_n z_n^{L-M+1} \end{bmatrix}_{n \times n} \quad (16)$$

The superscript T denotes matrix transpose. Note that the null space $\mathcal{N}(\mathbf{V}) = \mathbf{0} \in \mathbb{R}^M$ because $\mathbf{V}$ is a rectangular Vandermonde matrix with n rows and M columns such that $n > M$. Let $\mathbf{x}$ be a vector in $\mathbb{R}^M$ so that $\mathbf{V}^T \mathbf{\Lambda} \mathbf{V} \mathbf{x} = \mathbf{0}$. Consider the inner product $(\mathbf{x}, \mathbf{y}) := \sum_{k=1}^n x_k y_k$ for $\mathbf{x} = (x_1, \dots, x_n)^T$, $\mathbf{y} = (y_1, \dots, y_n)^T$, then

$$0 = (\mathbf{V}^T \mathbf{\Lambda} \mathbf{V} \mathbf{x}, \mathbf{x}) = (\mathbf{\Lambda} \mathbf{V} \mathbf{x}, \mathbf{V} \mathbf{x}) \geq \min_{i=1 \sim n} \lambda_i z_i^{L-M+1} (\mathbf{V} \mathbf{x}, \mathbf{V} \mathbf{x})$$

Since $\min_{i=1 \sim n} \lambda_i z_i^{L-M+1} > 0$, we have $\mathbf{V} \mathbf{x} = \mathbf{0}$. Hence $\mathbf{x} = \mathbf{0} \in \mathbb{R}^M$. This shows that the coefficient matrix in (8) is non-singular so the $[L/M]$ approximant exists for $L - M + 1 \geq 0$ and $1 \leq M < n$ if $z_i \neq 0$ for every i .

If $z_1 = 0$, $f(\xi)$ is a rational function with degree $n-1$ numerator and degree $n-1$ denominator. The characterization theorem of rational function (Theorem 1.4.4) in [8] implies that $[L/M](-\xi) = f(\xi)$ if $L \geq n-1$ and $M \geq n-1$. This takes care of $L - M + 1 \geq 1$ and $M \geq n-1$, represented by the triangular region marked by $\times$ in Table 1. For $L - M + 1 = 0$ and $M \leq n-1$, the coefficient matrix in (8) can be decomposed in the same fashion as in (14) with $\mathbf{\Lambda}$ replaced by

$$\mathbf{\Lambda} := \begin{bmatrix} \lambda_1 & 0 & 0 & \cdots & 0 \\ 0 & \lambda_2 & 0 & \cdots & 0 \\ 0 & 0 & \lambda_3 & \cdots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \cdots & \lambda_n \end{bmatrix}_{n \times n} \quad (17)$$

By an argument exactly the same as for the $z_1 \neq 0$ case, the existence of the $[L/M]$ Padé approximant for $L - M + 1 = 0$ and $M \leq n-1$ is proved; they correspond to the line formed by $\circ$ in

Table 1: Schematic description for case $n \geq 2$ and $z_1 = 0$

$M \setminus L$	0	1	2	$\dots$	$n-3$	$n-2$	$n-1$	n	$n+1$	$\dots$
0	•	•	•	$\dots$	•	•	•	•	•	$\dots$
1	○	•	•	$\dots$	•	•	•	•	•	$\dots$
2		○	•	$\dots$	•	•	•	•	•	$\dots$
3			○	$\dots$	•	•	•	•	•	$\dots$
$\vdots$				$\ddots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\vdots$	$\ddots$
$n-2$					○	•	•	•	•	$\dots$
$n-1$						○	$\times$	$\times$	$\times$	$\dots$
n							○	$\times$	$\times$	$\dots$
$n+1$								○	$\times$	$\dots$
$\vdots$									$\ddots$	$\ddots$

Table 1. For $L - M + 1 > 0$ and $M \leq n - 1$, the coefficient matrix of (8) is

$$\begin{bmatrix} \sum_{i=2}^n \lambda_i z_i^{L-M+1} & \sum_{i=2}^n \lambda_i z_i^{L-M+2} & \dots & \sum_{i=2}^n \lambda_i z_i^L \\ \sum_{i=2}^n \lambda_i z_i^{L-M+2} & \sum_{i=2}^n \lambda_i z_i^{L-M+3} & \dots & \sum_{i=2}^n \lambda_i z_i^{L+1} \\ \vdots & \vdots & \ddots & \vdots \\ \sum_{i=2}^n \lambda_i z_i^L & \sum_{i=2}^n \lambda_i z_i^{L+1} & \dots & \sum_{i=2}^n \lambda_i z_i^{L+M-1} \end{bmatrix} = \mathbf{V}^T \mathbf{\Lambda} \mathbf{V}, \quad (18)$$

where

$$\mathbf{V} := \begin{bmatrix} 1 & z_2 & z_2^2 & \dots & z_2^{M-1} \\ 1 & z_3 & z_3^2 & \dots & z_3^{M-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & z_M & z_M^2 & \dots & z_M^{M-1} \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 1 & z_n & z_n^2 & \dots & z_n^{M-1} \end{bmatrix}_{(n-1) \times M} \quad (19)$$

$$\mathbf{\Lambda} := \begin{bmatrix} \lambda_2 z_2^{L-M+1} & 0 & 0 & \dots & 0 \\ 0 & \lambda_3 z_3^{L-M+1} & 0 & \dots & 0 \\ 0 & 0 & \lambda_4 z_4^{L-M+1} & \dots & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & 0 & \dots & \lambda_n z_n^{L-M+1} \end{bmatrix}_{(n-1) \times (n-1)} \quad (20)$$

Since $\mathbf{V}$ is a Vandermonde matrix with $n - 1$ rows and M columns such that $n - 1 \geq M$ and $z_i \neq z_j$ for $i \neq j$, the null space $\mathcal{N}(\mathbf{V}) = \mathbf{0} \in \mathbb{R}^M$. Because all the entries in $\mathbf{\Lambda}$ are positive, the existence of the $[L/M]$ Padé approximant can be concluded in the same way as for the previous case. The corresponding region in the Padé table is marked in Table 1 by $\bullet$. $\square$

We apply this result to show the existence of accuracy-through-order Padé approximants for $G(\xi)$.

2.2 Standard Padé approximants for $G(\xi)$

Theorem 2.2. *The $[L/M]$ Padé approximant for $G(\xi)$ exists for all $L - M \geq 0$ and $M \geq 0$ such that $b_0 = 1$.*

Proof. For $L \geq 1$, let $[L - 1/M]_f$ denote the $[L - 1/M]$ Padé approximant of $f(\xi)$ such that $(L - 1) - M + 1 \geq 0$, $L \geq 1$ and $M \geq 0$. The proof follows immediately from the definition of G , Theorem 2.1 and definition of accuracy-through-order property (10):

$$G(\xi) - (-\xi)[L - 1/M]_f = -\xi(f(\xi) - [L - 1/M]_f) = O(\xi^{L+M+1}) \text{ for } |\xi| < 1.$$

That is, $(-\xi)[L - 1/M]_f$ is the $[L/M]$ Padé approximant of $G(\xi)$ for $L \geq 1$ and $L \geq M \geq 0$. If $L = 0$, then $M = 0$ because of the condition $L \geq M$. The $[0/0]$ approximant of G is 0 because $G(\xi) - 0 = \mu_0(-\xi) + \mu_1(-\xi)^2 + \dots = O(\xi^{0+0+1})$. The last equality is due to the positivity of μ_0 . $\square$

The algorithm presented in [2] for reconstructing spectral function from known measured data $\epsilon^*(\xi)$ for ξ not necessarily in $|\xi| < 1$. This can be justified by the two theorems listed below regarding properties of Padé approximants of Stieltjes series.

Theorem 2.3. *The poles of the Padé approximants of f , denoted by $[L/M]_f$, $L - M + 1 \geq 0$ and $M \geq 0$ are simple poles lying on $(-\infty, -1)$ with positive residues.*

Proof. For the case where the measure μ takes infinitely many different values, the proof can be found in [8], Theorem 5.2.1, p201. When μ takes $(n + 1) < \infty$ different values, the proof given in [8] needs to be modified as follows. The key step is to show the interlacing property of the zeros of $\triangle_M^{(J)}(x)$ defined in (21).

If $M = 0$, the Padé approximant has no pole. For $n = 0, 1$, the proof is the same as that for Theorem 2.1.

For $n \geq 2$, we first consider the case when $z_1 \neq 0$, i.e. no jump in values at $z = 0$ hence $f(\xi)$ is a rational function with numerator of degree $n - 1$ and denominator of degree n . Let $J := L - M \geq -1$, and $M \geq 1$. Following the proof of Theorem 5.2.1 in [8], we define

$\Delta_0^{(J)}(x) = 1$; for $M = 1, 2, 3 \dots$, define

$$\Delta_M^{(J)}(x) := \begin{vmatrix} \mu_{1+J} + x\mu_{2+J} & \mu_{2+J} + x\mu_{3+J} & \cdots & \mu_{M+J} + x\mu_{M+J+1} \\ \mu_{2+J} + x\mu_{3+J} & \mu_{3+J} + x\mu_{4+J} & \cdots & \mu_{M+J+1} + x\mu_{M+J+2} \\ \vdots & \vdots & \ddots & \vdots \\ \mu_{M+J} + x\mu_{M+J+1} & \mu_{M+J+1} + x\mu_{M+J+2} & \cdots & \mu_{2M+J-1} + x\mu_{2M+J} \end{vmatrix} \quad (21)$$

Note that $\Delta_M^{(J)}(x) = Q^{[L/M]}(-x)$ by applying (3.2) on p.16 of [8] to (13) and that $\Delta_M^{(J)}(0) = 0$ for $M > n$ and $J \geq -1$. Hence the zeros of $\Delta_M^{(J)}$ correspond to the poles of Padé approximant $[L/M]$. Unlike the proof for $n = \infty$ in [8], Theorem 5.2.1, we only need to consider $J \geq -1$ and $1 \leq M < n$ because for $J \geq -1$ and $M \geq n$, the Padé approximant for $f(\xi)$ is itself the poles of $f(\xi)$ with finite n are apparently all simple and located in $(-\infty, -1)$. Next, we need to show that the functions $\Delta_1^{(J)}(x), \dots, \Delta_{n-1}^{(J)}(x)$ have the property such that if $\Delta_M^{(J)}(x) = 0$ for some x , then $\Delta_{M-1}^{(J)}(x)$ and $\Delta_{M+1}^{(J)}(x)$ have opposite signs for all $1 \leq M \leq n-1$. This can be shown by using Sylvester's identity and Frobenius identity in exactly the same way as in the proof in [8], Theorem 5.2.1. In order to show the interlacing property of the zeros of $\Delta_M^{(J)}(x)$, we need to prove the following properties

$$\left. \begin{aligned} \Delta_M^{(J)}(0) &> 0 \\ \Delta_M^{(J)}(-\infty) &= (-1)^M \cdot \infty \end{aligned} \right\} \text{ for } M = 1, \dots, n \quad (22)$$

by first noting that $\Delta_M^{(J)}(0)$ is the determinant of the matrix $\mathbf{V}^T \mathbf{\Lambda} \mathbf{V}$ in (14) and

$$(\mathbf{V}^T \mathbf{\Lambda} \mathbf{V} \mathbf{y}, \mathbf{y}) = (\mathbf{\Lambda} \mathbf{V} \mathbf{y}, \mathbf{V} \mathbf{y}) \geq \min_{j=1, \dots, n} \lambda_j z_j^{L-M+1} (\mathbf{V} \mathbf{y}, \mathbf{V} \mathbf{y}) > 0, \text{ for } \mathbf{y} \in \mathbb{R}^M \setminus \{\mathbf{0}\}$$

So $\det(\mathbf{V}^T \mathbf{\Lambda} \mathbf{V}) > 0$, hence $\Delta_M^{(J)}(0) > 0$ for $M = 0, 1, \dots, n$. For $\Delta_M^{(J)}(-\infty)$, we decompose the matrix corresponding to $\Delta_M^{(J)}(x)$ as

$$\begin{bmatrix} \mu_{1+J} + x\mu_{2+J} & \mu_{2+J} + x\mu_{3+J} & \cdots & \mu_{M+J} + x\mu_{M+J+1} \\ \mu_{2+J} + x\mu_{3+J} & \mu_{3+J} + x\mu_{4+J} & \cdots & \mu_{M+J+1} + x\mu_{M+J+2} \\ \vdots & \vdots & \ddots & \vdots \\ \mu_{M+J} + x\mu_{M+J+1} & \mu_{M+J+1} + x\mu_{M+J+2} & \cdots & \mu_{2M+J-1} + x\mu_{2M+J} \end{bmatrix} = \mathbf{V}^T \mathbf{\Lambda}_x \mathbf{V}$$

with

$$\mathbf{\Lambda}_x := \begin{bmatrix} \lambda_1 z_1^{L-M+1} (1 + z_1 x) & 0 & \cdots & 0 \\ 0 & \lambda_2 z_2^{L-M+1} (1 + z_2 x) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n z_n^{L-M+1} (1 + z_n x) \end{bmatrix}_{n \times n}$$

Similar as before, for a fixed x , there exists A and B such that

$$A(\mathbf{V}\mathbf{y}, \mathbf{V}\mathbf{y}) \geq (\mathbf{V}^T \mathbf{\Lambda}_x \mathbf{V}\mathbf{y}, \mathbf{y}) = (\mathbf{\Lambda}_x \mathbf{V}\mathbf{y}, \mathbf{V}\mathbf{y}) \geq B(\mathbf{V}\mathbf{y}, \mathbf{V}\mathbf{y}), \text{ for } \mathbf{y} \in \mathbb{R}^M \setminus \{\mathbf{0}\}$$

where

$$A := \max_{j=1, \dots, n} \lambda_j z_j^{L-M+1} (1 + z_j x), \quad B := \min_{j=1, \dots, n} \lambda_j z_j^{L-M+1} (1 + z_j x)$$

As $x \rightarrow -\infty$, we have $A \rightarrow -\infty$ and $B \rightarrow -\infty$. Hence $\Delta_M^{(J)}(x) = \det(\mathbf{V}^T \mathbf{\Lambda}_x \mathbf{V}) \rightarrow (-\infty)^M$ as $x \rightarrow -\infty$. This completes the proof of (22) for $z_1 \neq 0$.

For $z_1 = 0$, we first consider that case when $L - M + 1 = 0$ (i.e. $J = -1$) and $n - 1 \geq M \geq 1$. We have $\Delta_M^{(J)}(0) = \det(\mathbf{V}^T \mathbf{\Lambda} \mathbf{V})$ with $\mathbf{\Lambda}$ defined in (17). So $\Delta_M^{(J)}(0) > 0$ for the same reason as in the $z_1 \neq 0$ case. As for $\Delta_M^{(J)}(x)$ for fixed x , we have $\Delta_1^{(J)}(x) = \lambda_1 + \sum_{i=2}^n \lambda_i (1 + x z_i) \rightarrow -\infty$ as $x \rightarrow -\infty$ and for $M \geq 2$

$$\Delta_M^{(J)}(x) = \det \left(\begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{bmatrix}_{M \times M} + \mathbf{V}^T \mathbf{\Lambda}_x \mathbf{V} \right)$$

with $\mathbf{V}$ being the one defined in (19) and

$$\mathbf{\Lambda}_x := \begin{bmatrix} \lambda_2(1 + z_2 x) & 0 & \cdots & 0 \\ 0 & \lambda_3(1 + z_3 x) & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & \lambda_n(1 + z_n x) \end{bmatrix}_{(n-1) \times (n-1)} \quad (23)$$

Therefore, for any $\mathbf{y} = (y_1, \dots, y_M)^T \neq \mathbf{0}$, we have

$$\lambda_1 y_1^2 + A(\mathbf{V}\mathbf{y}, \mathbf{V}\mathbf{y}) \geq \left(\left(\begin{bmatrix} \lambda_1 & 0 & \cdots & 0 \\ \vdots & \vdots & \ddots & \vdots \\ 0 & 0 & \cdots & 0 \end{bmatrix}_{M \times M} + \mathbf{V}^T \mathbf{\Lambda}_x \mathbf{V} \right) \mathbf{y}, \mathbf{y} \right) \geq \lambda_1 y_1^2 + B(\mathbf{V}\mathbf{y}, \mathbf{V}\mathbf{y})$$

with $A := \max_{i=2, \dots, n} \lambda_i (1 + z_i x)$ and $B := \min_{i=2, \dots, n} \lambda_i (1 + z_i x)$; hence $\Delta_M^{(J)}(x) \rightarrow (-\infty)^M$ as $x \rightarrow -\infty$. For $L - M + 1 \geq 1$, $\Delta_M^{(J)}(0) = \det(\mathbf{V}^T \mathbf{\Lambda} \mathbf{V})$ and $\Delta_M^{(J)}(x) = \det(\mathbf{V}^T \mathbf{\Lambda}_x \mathbf{V})$ with $\mathbf{V}$ defined in (19), $\mathbf{\Lambda}$ defined in (20) and $\mathbf{\Lambda}_x$ defined in (23) and the same argument can be applied for proving (22). The rest of the proof is the same as that in [8], Theorem 5.2.1. $\square$

This theorem implies the following theorem:

Theorem 2.4 ([8], Theorem 5.4.4, p225). *Any sequence of $[L_k/M_k]$ Padé approximants, $L_k - M_k + 1 \geq 0$, of a Stieltjes series $f(\xi)$ convergent in $|\xi| < 1$ converges uniformly to $f(\xi)$ in $\mathcal{D}^+(\Delta)$, where $\mathcal{D}^+(\Delta)$ is the set of all points in $|\xi| < R$ which are at least Δ distance away from the cut $(-\infty, -1]$ for any positive numbers R and Δ .*

In other words, even though the Padé approximants are constructed from the Stieltjes series convergent only in $|\xi| < 1$, their validity of approximating the Stieltjes function $f(\xi)$, which is analytic everywhere outside $(-\infty, -1]$, extends far beyond $|\xi| < 1$.

In [4], Cherkaev et. al. define S_N equivalence of dielectric composites with isotropic constituents. According to their definition, two microstructures are S_N -equivalent if the the first N moments (μ_0 to μ_{n-1}) of the spectral measures $\mu_{ik}^{(1)}$ and $\mu_{ik}^{(2)}$ in the IRF (1) are identical. In the following, we show that Padé approximants provide a characterization of S_N -equivalent structures in the following sense:

Theorem 2.5. Let $G_1(\xi)$ and $G_2(\xi)$ be two Stieltjes functions defined as in (5) by the effective complex dielectric constant of Structure-1 and Structure-2, respectively. Denote the corresponding $[L/M]$ Padé approximant by $[L/M]_{G_1}$ and $[L/M]_{G_2}$. Then Structure-1 and Structure-2 are S_N equivalent, $N \geq 1$, if and only if $[L/M]_{G_1} = [L/M]_{G_2}$ for all $L + M \leq N$ such that $L \geq M \geq 0$.

Proof. Since the coefficients of $(-\xi)^{n+1}$, $n \geq 0$, in the power series expansions of G_1 and G_2 (7) are given by the n^{th} moment of the corresponding measure, S_N -equivalence implies that the power series expansions near $\xi = 0$ of G_1 and G_2 have the same first $N + 1$ terms (the constant term in the series is 0 for both G_1 and G_2). Since the $[L/M]$ Padé approximant is completely determined by coefficients up to $(-\xi)^{L+M}$, clearly $[L/M]_{G_1} = [L/M]_{G_2}$ if $L + M \leq N$ and if they both exist. By Theorem 3.1, we conclude that $[L/M]_{G_1} = [L/M]_{G_2}$ for $L + M \leq N$ and $L \geq M \geq 0$.

Conversely, suppose $[L/M]_{G_1}(-\xi) = [L/M]_{G_2}(-\xi)$ for $L + M = N$. Due to Theorem 2.3 all the poles of Padé approximants for Stieltjes series convergent in $|\xi| < 1$ lie on the real axis in the interval $-\infty < \xi < -1$, the power series expansion of this rational function

$$[L/M]_{G_1}(-\xi) = [L/M]_{G_2}(-\xi) = \sum_{n=0}^{\infty} c_n^{[L/M]} (-\xi)^n.$$

is valid for $|\xi| < 1$. By accuracy-through-order property of the Padé approximants, $c_n^{[L/M]} = \mu_{n-1}^{(1)}$ and $c_n^{[L/M]} = \mu_{n-1}^{(2)}$ for $n = 1 \sim N$ with $\mu_n^{(1)}$ and $\mu_n^{(2)}$ being the coefficients of G_1 and G_2 , respectively, as defined in (7). Hence the theorem is proved. $\square$

3 Existence of nonstandard Padé approximants for $F(s)$

The main result of this paper is to show the existence of nonstandard Padé approximants in (2).

Theorem 3.1. *For any Stieltjes function*

$$F(s) = \int_0^1 \frac{1}{s-z} d\mu(z) \quad (24)$$

with non-decreasing spectral function $\mu(z)$ taking either infinitely or finitely many different values for $z \in [0, 1]$, there exists nonstandard Padé approximant of the form

$$F(s) \simeq \frac{a(s)}{b(s)} = \frac{a_0 + a_1s + a_2s^2 + \cdots + a_{M-1}s^{M-1}}{b_0 + b_1s + b_2s^2 + \cdots + b_Ms^M}, \quad b_1 \neq 0, M \geq 1. \quad (25)$$

Proof. We first consider the case in which $\mu(z)$ takes infinitely many different values on $[0, 1]$. Let $L = M - 1$ and $M \geq 1$. According to Theorem 2.1 and Theorem 2.3, the $[M - 1/M]$ Padé approximant of $f(\xi)$ exists with simple poles lying on $\xi \in (-\infty, -1)$ with positive residues. The interlacing-of-zero property of $\triangle_M^{(J)}(x)$ guarantees the existence of M different poles of $[M - 1/M]$. By Theorem 2.3, it can be further expressed in terms of the partial fraction decomposition

$$[M - 1/M]_f = \frac{P(-\xi)}{Q(-\xi)} = \sum_{i=1}^M \frac{\lambda_i}{\xi - p_i}, \quad p_i \in (-\infty, -1), \quad p_i \neq p_j \quad \text{for } i \neq j, \quad \lambda_i > 0. \quad (26)$$

This induces the $[M/M]$ approximant of $G(\xi)$:

$$[M/M]_G = -\xi \cdot [L/M]_f = \sum_{i=1}^M \frac{-\xi \cdot \lambda_i}{\xi - p_i}, \quad p_i \in (-\infty, -1), \quad \lambda_i > 0. \quad (27)$$

We rewrite the expression above in terms of variable $s := \frac{-1}{\xi} \in [0, 1)$

$$[M/M]_G = \sum_{i=1}^M \frac{-\xi \cdot \lambda_i}{\xi - p_i} = \sum_{i=1}^M \frac{-\lambda_i/p_i}{s - (-1/p_i)}. \quad (28)$$

Noting $\frac{-1}{p_i} \in [0, 1)$ and $F(s) = G(\xi)$, the partial fraction representation above implies there exists a rational function approximation of $F(s)$ with the following form

$$F(s) \simeq \frac{a(s)}{b(s)} = \frac{a_0 + a_1s + a_2s^2 + \cdots + a_{M-1}s^{M-1}}{b_0 + b_1s + b_2s^2 + \cdots + b_Ms^M}, \quad M \geq 1$$

where $b_1 \neq 0$. This is because the coefficient of s in the denominator of (28) is $\sum_{i=1}^M \prod_{j \neq i} \frac{1}{p_j}$, which is non-zero for $M \geq 2$. For $M = 1$, this is obviously true.

Next, we consider the case for μ taking $(n + 1)$ different values, $1 \leq n < \infty$ with jumps at $1 > z_1, \dots, z_n \geq 0$ of magnitude $\lambda_1, \dots, \lambda_n > 0$. Then

$$F(s) = \sum_{i=1}^n \frac{\lambda_i}{s - z_i} \quad (29)$$

For $M \geq n$, the $[M - 1/M]$ Padé approximant of $F(s)$ is itself and apparently (29) has the form (25). For $M < n$, the proof is the same as in the case where μ takes infinitely many different values because of the validity of Theorem 2.1 and Theorem 2.3 for this case. $\square$

4 Conclusion and discussion

In this paper, we prove that a Stieltjes function of the form (24) can be approximated by $[M - 1/M]$ non-standard Padé approximant (25). Most of the work in this paper is dedicated to the case where the spectral measure μ takes finitely many different values in $[0, 1]$ because the proof for μ that takes infinitely many different values in $[0, 1]$ can be obtained by results that are already available in the literature, as indicated in the theorems presented in this paper. Most results in the literature concern about the convergence of $[L/M]$ Padé approximants to a Stieltjes function as L and/or $M \rightarrow \infty$. Naturally, they exclude those functions whose spectral function μ takes finitely many different values due to the fact that these functions are rational functions and $[L/M]$ Padé approximants for a rational function is the function itself when L and M are large enough; hence convergence is guaranteed. The motivation for proving the main result presented in Theorem 3.1 is due to the concerns on its application on inverse homogenization as described in the introduction. It's well known that for any rank n laminate microstructure, $F(s)$ has finite number of poles, i.e. its spectral measure takes finite number of different values. At the same time, it is also known that the checkerboard microstructure has a spectral function which takes infinitely many different values and has a jump at $z = 0$. In other words, given data on effective dielectric constant of a composite, we only know that the corresponding $F(s)$ function can be represented as a Stieltjes integral but we do not know *a priori* how many different values the spectral function takes because the microstructure itself is the unknown for the inverse problem. From the inverse problem point of view, the existence of the nonstandard Padé approximants is as important as the convergence property. As mentioned in the introduction, in [2], the spectral measure was reconstructed with very good precision by using data of effective dielectric constants at different frequencies. It is done by recovering the coefficients $b_0, b_2, \dots, b_M, a_0, \dots, a_M$ by reformulating the problem as minimization problem with constraints on the location of poles s_n and residues A_n in (4). The results in this paper also justify these constraints except $\sum_n A_n < 1$ by identifying A_n with $-\frac{\lambda_n}{p_n}$ and s_n with $-1/p_n$ in (28). The constraint $\sum_n A_n < 1$ is justified by noticing that $\sum_i \frac{-\lambda_i}{p_i}$ is the constant term of the Maclaurin series of $[M/M - 1]_f$ in (26) and it has to be equal to the constant term of the Maclaurin series of f , which is μ_0 in (6). $\mu_0 < 1$ because it is the volume fraction of one of the constituent materials.

Acknowledgments The author would like to thank E. Cherkaev and D. Zhang for bringing into her attention the problem regarding existence of nonstandard Padé approximants for IRF. This research is partially sponsored by the ARRA-NSF Grant DMS-0920852.

References

- [1] K. Golden and G. Papanicolaou , Bounds on effective parameters of heterogeneous media by analytic continuation, *Commun. Math. Phys.* **90** (1983) 473–491.
- [2] D-L. Zhang and E. Cherkaev, *Journal of Computational Physics* **228** (2009) 5390–5409.
- [3] E. Cherkaev and M-J. Yvonne Ou, Dehomogenization: reconstruction of moments of the spectral measure of the composite, *Inverse Problems* **24** (2008) 065008.
- [4] E. Cherkaev and D-L. Zhang, De-homogenization: equivalent classes of microgeometries of composites and moments of the spectral function reconstructed from effective complex permittivity *Preprint* (2009).
- [5] Christian Engström, Bounds on the effective tensor and the structural parameters for anisotropic two-phase composite material, *J. Phys. D: Appl. Phys.* **38** (2005) 3695–3702 .
- [6] S. Tokarzewski and J. Telega, A contribution to the bounds on real effective moduli of two-phase composite materials , *Mathematical Models and Methods in Applied Sciences.* **7(6)** (1997) 769–789.
- [7] S. Tokarzewski and J. Telega, Bounds on effective moduli by analytical continuation of the Stieltjes function expanded at zero and infinity, *Zeitschrift für angewandte Mathematik und Physik* **49** (1998) 137–155.
- [8] George A. Baker Jr and Peter Graves-Morris, *Padé Approximants* (Cambridge University Press; Second edition, 1996)
- [9] Elena Cherkaev, Inverse homogenization for evaluation of effective properties of a mixture , *Inverse Problems* **17** (2001) 1203–1218.